

# Reviewing working memory training gains in healthy older adults: A meta-analytic review of transfer for cognitive outcomes

Ana C. Teixeira-Santos<sup>a</sup>, Célia S. Moreira<sup>b,c</sup>, Rosana Magalhães<sup>d,e</sup>, Carina Magalhães<sup>b</sup>, Diana R. Pereira<sup>a</sup>, Jorge Leite<sup>a,f,g</sup>, Sandra Carvalho<sup>a,f</sup>, Adriana Sampaio<sup>a</sup>

<sup>a</sup>Psychological Neuroscience Laboratory–CIPsi, School of Psychology, University of Minho, Campus de Gualtar, Braga, Portugal

<sup>b</sup>CIPsi, School of Psychology, University of Minho

<sup>c</sup>Department of Mathematics, University of Porto (FCUP, CMUP)

<sup>d</sup>Serviço de Psiquiatria/Psicologia do Hospital Cuf Porto

<sup>e</sup>Universidade Católica Portuguesa, CEDH – Centro de Investigação para o Desenvolvimento Humano, HNL – Human Neurobehavioral Laboratory, Porto – Portugal

<sup>f</sup>Spaulding Neuromodulation Center, Department of Physical Medicine & Rehabilitation, Spaulding Rehabilitation Hospital and Massachusetts General Hospital, Harvard Medical School, Boston, USA

<sup>g</sup>Portugalense Institute for Human Development (INPP), Universidade Portugalense, Porto, Portugal

## ARTICLE INFO

### Keywords:

Working memory training  
Cognitive plasticity  
Training transfer  
Healthy older adults  
Healthy ageing.

## ABSTRACT

The objective of this meta-analytic review was to systematically assess the effects of working memory training on healthy older adults. We identified 552 entries, of which 27 experiments met our inclusion criteria. The final database included 1130 participants. Near- and far-transfer effects were analysed with measures of short-term memory, working memory, and reasoning. Small significant and long-lasting transfer gains were observed in working memory tasks. Effects on reasoning was very small and only marginally significant. The effects of working memory training on both near and far transfer in older adults were moderated by the type of training tasks; the adopted outcome measures; the training duration; and the total number of training hours. In this review, we provide an updated review of the literature in the field by carrying out a robust multi-level meta-analysis focused exclusively on working memory training in healthy older adults. Recommendations for future research are suggested.

## 1. Introduction

Ageing of the world population is a major public health concern that has captured the attention of the general public. Overall, more than 962 million people were over the age of 60 in 2017. It is estimated that this number will more than double to 2.1 billion people by the year 2050 (United Nations, Department of Economic and Social Affairs, 2017). Specifically, it is estimated that the population of people over the age of 80 will triple by the year 2050, increasing from 137 million to 425 million (United Nations, Department of Economic and Social Affairs, 2017). Therefore, much effort has been made to promote optimal ageing to avoid both declines in cognitive functioning and dependence on others, which are factors associated with ageing. Specifically, much has been done to try to reverse age-related cognitive decline and prevent or delay pathological cognitive disorders. This movement represents a significant attempt to improve the quality of life of older adults and to relieve the burden on medical care systems that has

resulted from a substantial increase in the elderly population. Efforts to address the issue include non-pharmacological interventions, such as the Advanced Cognitive Training for Independent and Vital Elderly (ACTIVE) trial (Ball et al., 2002, 2002; Rebok et al., 2014). The promising findings in the field encouraged researchers to further investigate the benefits of cognitive training in older people.

Different cognitive training approaches are reported in the literature (Jolles and Crone, 2012). They can be classified into two major categories: “strategy-based training” and “process-based approaches”. *Strategy-based training* consists of the development of specific adaptations and strategies, such as mnemonics, which can be used to ameliorate daily struggles (Lustig et al., 2009), whereas process-based approaches focus on the training of specific cognitive abilities (Clare and Woods, 2004). More specifically, *core process-based training* focuses on training central mechanisms with the purpose of producing more substantial effects in functions that depend upon this central processor and that share a common neural substrate (Morrison and Chein, 2011).

Notably, working memory training (WMT) has emerged as a proxy for improving cognitive functions (Neely and Nyberg, 2015).

Working memory (WM) refers to the components responsible for maintain temporally a limited amount of information in an available state to allow the processing of ongoing information (Cowan, 2017). WM performance declines markedly with ageing, and this has been associated with abnormalities on the frontoparietal networks involved in WM, as well as neuromodulatory (dopamine) and neuroanatomical alterations (Bäckman et al., 2017, 2010; Lubitz et al., 2017; Park and Reuter-Lorenz, 2009; Raz, 2005; Rottschy et al., 2012; Salthouse, 1990). This reduction in WM capacity in older adults, along with a decrease in processing speed, seem to underlie age-related cognitive decline (Braver and West, 2008), primarily because WM is associated with higher-order cognitive functions (Unsworth et al., 2005), including reasoning (Shakeel and Goghari, 2017), reading (Just and Carpenter, 1992), prospective memory (Bisiacchi et al., 2008), processing speed (Diamond et al., 1999), attention (West, 1999), perceptual organization (Ko et al., 2014), and general language (Kemper et al., 2004). Therefore, given the decrease in WM performance with ageing and its putative role in higher-order cognitive functions, WMT has been studied extensively to enhance cognition in older adults, and positive effects of WMT on both cognition and neural plasticity have been found (Constantinidis and Klingberg, 2016; Karbach and Verhaeghen, 2014).

Experimental studies of WMT typically include an experimental group, whose members participate in a WMT, and a control group. The control group can be a no-contact control group (passive control group) or an active control group that completes a non-related activity or a low-level WMT. Participants in active control group are exposed to a training setting (i.e., number of sessions, contact with the experimenter, a style of intervention) that is similar to that of the experimental group, but they are not exposed to the experimental WM condition. This design with active control condition allows the researcher to control for effects that may result from social contact during the experiment or a participant's expectations. However, participants from both groups (passive and active control groups) undergo the same testing before and after the intervention as the participants of the experimental groups.

There is abundant literature on WMT (see app. table 4). They may include computerized tasks and can be visual, auditory or both visual and auditory. Trained tasks usually consist of complex or simple span tasks or updating tasks. In complex span tasks, participants must recall a sequence of stimuli, which is interleaved with a concurrent activity. In simple span tasks, participants must remember the sequence of stimuli in forward (fwd) or backward (bwd) order. Updating includes tasks in which participants hold specific content in memory, continually up-dating the information to be remembered and dropping information that is no longer needed. Training is usually adaptive, i.e., the task difficulty adjusts based on the individual's performance (von Bastian and Eschen, 2016).

Several studies have been designed to study the effects of WMT by comparing the pre- and posttest results of experimental and control groups immediately after training (posttest) and at a delayed post-training assessment (follow-up). Additionally, studies have investigated the transfer effects, i.e., whether training gains can be generalized to other tasks involving different cognitive abilities (e.g., Borella et al., 2010) such as fluid intelligence (Beatty et al., 2015). Although there are no clear criteria to define transfer distance, most authors locate the generalization of the effects along a continuum of near to far transfer (Noack et al., 2009). Near transfer consists of an improvement on tasks that are like the trained task and that share the same mechanisms or components, while far transfer represents an improvement on tasks that measure abilities that are not like the abilities trained. Near-transfer effects are commonly observed (Borella et al., 2010; Li et al., 2008), although this is not always the case (Dahlin et al., 2008a). Results re-garding far transfer are controversial with limited or no evidence (Borella et al., 2013).

Previous narrative and systematic reviews have debated the

potentialities and controversies of WMT (Constantinidis and Klingberg, 2016; Karbach and Verhaeghen, 2014; Lampit et al., 2014; Melby-Lervåg and Hulme, 2016, 2013; Morrison and Chein, 2011; Schwaighofer et al., 2015; von Bastian and Oberauer, 2013; Weicker et al., 2016), yet the results are inconclusive (see app. Table 1). Therefore, the current meta-analysis aims to contribute to this debate by examining the generalization of training effects to non-trained tasks (near and far transfer) (aim 1) and the maintenance of the effects over time (i.e., at follow-up) (aim 2) by using a meta-analysis approach that is different from the ones used in previous reviews.

Additionally, previous meta-analyses (Karbach and Verhaeghen, 2014; Melby-Lervåg et al., 2016) and experimental studies (e.g., Bürki et al., 2014; Stepankova et al., 2014; Zinke et al., 2014) have suggested that variables such as type of control group (Melby-Lervåg et al., 2016), age (Bürki et al., 2014; Richmond et al., 2011; von Bastian et al., 2013; Zajac-Lamparska and Trempala, 2016; for works considering only older adults, see: Borella et al., 2017a; Zinke et al., 2014), education (Borella et al., 2017a), general cognitive ability (Borella et al., 2017a), baseline performance (Zinke et al., 2014, 2011) and training dosage (Bürki et al., 2014; Lilienthal et al., 2013; Stepankova et al., 2014) might moderate training gains and transfer effects. For instance, in relation to the type of control group, a meta-analysis from Melby-Lervåg et al. (2016) re-ported that the type of control group predicted transfer effects. In particular, studies showed more significant effects when using a passive control group than when using an active control group. However, other meta-analytical studies (Karbach and Verhaeghen, 2014; Weicker et al., 2016) did not find influence of type of control group (active or passive) in transfer effects. Regarding the age, an experimental study performed by Borella et al. (2014) found transfer effects of a visuospatial WMT for measures of STM, WM, inhibition, processing speed, and reasoning only in young-old adults but not in old-old adults. In accordance, together with an age-related difference in the transfer effects, Borella et al. (2017a) also documented the role of age as an important moderator of the effects in WMT, although the results varied according to the type of transfer task. In addition, Zinke et al. (2014) evidenced that old-old participants had less gains than young-old participants, except for fluid intelligence in which the reverse pattern was verified.

Borella et al. (2017a, 2017b) have also shown that vocabulary and baseline performance influenced WMT. In this study participants with higher vocabulary scores and poor pretest performance benefited more from training, although this pattern was not the same in all outcomes (e.g., in fwd digit span, lower vocabulary score was related to more benefit in training). Moreover, participants with low levels of baseline performance in WM tasks were likely to benefit more from WMT (Zinke et al., 2014, 2011). Related to session length/duration, Jaeggi et al (2008) documented a significant growth in far transfer throughout the sessions (from 8 to 19 sessions). Other researchers showed that a group which trained for 20-day outperformed a 10-day training group in a visuospatial measure (Stepankova et al., 2014), while a small positive significant moderator effect for small training dose in comparison to large training dose was observed in a meta-analysis (Melby-Lervåg et al., 2016).

Taken together, in the current study, we verified if the variables as type of control (active/passive), mean age of participants, total number of training hours, number of training sessions, training length in weeks, training type (single training - complex span, simple span, updating, or mixed training: more than one type of WM task), years of formal education, general cognitive ability (operationalized by vocabulary score), and baseline performance would moderate the training effect (aim 3). In addition, we also verified if the type of the outcome adopted (e.g., Cattell; Raven Advanced Progressive Matrix - RAPM; complex span) would moderate the transfer effect.

Previous meta-analytical work merged the results of different age groups (Mansur-Alves and Silva, 2017; Melby-Lervåg et al., 2016; Melby-Lervåg and Hulme, 2013) or did not include older adults (Au et al., 2015). This review focuses on only older adults, as WM is

markedly affected by ageing (Salthouse, 2000), and WMT is proposed as an innovative approach to counteract age-related cognitive declines (Constantinidis and Klingberg, 2016; Karbach and Verhaeghen, 2014). While merging different ages and conditions may yield sample heterogeneity, this practice can pose some problems for the internal and external validity of the findings (Rothwell, 2006). Additionally, to better isolate the effects of WMT, this meta-analysis addresses the specificity of the training delivered to the experimental groups by including studies whose experimental groups participated in trainings focused exclusively on WM and excluding studies whose experimental groups participated in trainings targeting cognitive functions other than WM. We also excluded papers whose active control groups participated in a non-adaptive WMT that remained always in a lower level of WMT (Brehmer et al., 2011; Chan et al., 2015; Loosli et al., 2016; Shing et al., 2012; Simon et al., 2018; Wayne et al., 2016), specific examples include: comparing an adaptive WMT with a WMT whose load (e.g.,  $N = 2$  or  $N = 3$ ) is held constant throughout the training (Brehmer et al., 2011; Chan et al., 2015; Wayne et al., 2016); training both experimental and control groups with a recent-probe and an  $n$ -back task, with the experimental group receiving trials with higher proactive interference when compared to the control group (Loosli et al., 2016); the participants performed a numerical memory updating task, however different groups were exposed to distinct rates of stimuli presentation (750 ms, 1500 ms or 3000 ms) (Shing et al., 2012). Considering that our aim was to contrast WMT with a placebo training not related to WM (e.g., questionnaire, quiz, visual search) or a non-training condition, in the present review, the above-mentioned studies were not included in the analysis. The rationale behind this is the fact that even a low-level of WM performance activates similar brain areas as high-level of WM processing (Braver et al., 1997; Kawagoe et al., 2015; Ragland et al., 2002). Since we do not have enough information to determine a sub-optimal dosage of WMT that would work solely as placebo (Huitfeldt et al., 2001), comparing different loads of WMT could lead to less interpretable data as these WM tasks might produce similar effects. As a consequence, we would not be able to isolate gains that are due to WMT (ICH Harmonised Tripartite Guideline, 2000). In fact, as suggested by Brehmer and colleagues (2011), both adaptive WMT and training at low WM load might lead to neural changes. Additionally, although many researchers classify executive function tasks as WM we did not include training of executive functions, such as Stroop interference, verbal fluency or task switching. As claimed by Oberauer et al. (2018) in the Benchmarks for Models of Short Term and Working Memory, executive functions are framed under specific theories and models that are different from the WM literature. Furthermore, similar to previous meta-analysis (Karbach and Verhaeghen, 2014), we focused on healthy older adults, which represents the majority of the aging population, grounded on the basis of maintenance or enhancement of cognition as a pre-ventive measure, instead of rehabilitation in non-normative aging as a remedial measure (Tkatch et al., 2016).

Regarding the methods carried out in this meta-analysis, we employed robust analytical methods to address multiple outcomes (Moeyaert et al., 2017) rather than use the average of the outcomes (e.g., Karbach and Verhaeghen, 2014; Melby-Lervåg and Hulme, 2016). Robust approaches to address multiple outcomes and treatments are critical as they give unbiased parameter estimates, while the average method may bias the estimates of the standard errors (Moeyaert et al., 2017; Morris, 2008). Finally, a sensitivity analysis was performed to address the lack of data on correlations between pre- and post-training measures. These correlational data are necessary to calculate the variance of the effect size of intervention gains, which was not considered in previous meta-analyses (Mansur-Alves and Silva, 2017; Melby-Lervåg and Hulme, 2016, 2013; Schwaighofer et al., 2015). Finally, a descriptive analysis of the risk of bias was provided following the Cochrane recommendations (Higgins and Altman, 2008). Overall, considering these methodological issues and the fact that new papers have been published since the publication of the most recent meta-analysis,

the current study offers an integrated and updated overview of WMT gains in healthy older adults in accordance with the Cochrane recommendations (Higgins and Green, 2008) that highlight the need to update reviews every two years.

## 2. Methods

We performed a systematic review following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA) (Moher et al., 2009).

### 2.1. Data sources and eligibility criteria

Five databases (Scopus, Pubmed, PsychINFO, Science Direct, and Scielo; the last was accessed through Web of Science) were searched on January 16, 2019. There were no time or language restrictions. The search terms used were “training”, “working memory”, and “older adult”. The combinations of descriptors can be found in the supplementary material (see table A). Additionally, reference lists from six major reviews and one book chapter in the field were also searched (Karbach and Verhaeghen, 2014; Melby-Lervåg et al., 2016; Melby-Lervåg and Hulme, 2016, 2013; Morrison and Chein, 2011; Noack et al., 2009; Shipstead et al., 2012). App Table 2 includes the inclusion/exclusion criteria, and app. Fig. A1 provides a schematic representation of the steps adopted in the literature search. When there were insufficient data to proceed with effect size estimations, an email was sent to the authors requesting the relevant information. In cases in which a reply from the authors was not possible, we limited the inclusion of the study to the data provided.

### 2.2. Data extraction

Two authors selected articles based on the titles and abstracts, and duplicate records were manually removed. After the exclusion of irrelevant articles, two authors independently performed a full-text analysis to assess the eligibility of the articles for inclusion in the review. There was moderate Fleiss' Kappa inter-rater reliability agreement between investigators in the full-text screening, including both included and excluded studies ( $k = 0.5$ ) (Landis and Koch, 1977). Two reviewers independently assessed the risk of bias using the Cochrane Collaboration's risk of bias tool (Higgins and Altman, 2008). Studies were classified as “high risk”, “low risk” or “unclear” in the following domains: randomization, concealment of allocation, blinding of participants, personnel and outcome assessment, attrition, and reporting bias. At any stage, disagreements between reviewers were solved with discussions or in consultation with a third reviewer. Statistical analyses were conducted using the R packages “metafor” (Viechtbauer, 2010), “forest-plot” (Gordon and Lumley, 2016), “clubSandwich”, (Pustejovsky, 2017), and “metaLik” (Guolo and Varin, 2012) from R statistical environment (RStudio, version 3.5.2, R Core Team, 2018).

Two reviewers independently recorded the following information from each full-text article: scores, standard deviations of pre- and post-treatment assessments, number of participants per group, types of outcomes, predictor variables, and dropout rates. Completion rates (i.e., the percentage of participants who completed training programmes) were calculated for each group. When a trial had two control groups, an active and a passive, we analysed data from the active group, as it is suggested that this approach allows better control of expectancy effects, such as the Hawthorne effect (Wickstrom and Bendix, 2000). One exception was the study of Weicker et al. (2018), from which the passive control group was selected instead of the active control group, as the latter performed a fixed low-level WM task (see app Table 2 for exclusion criteria).

To assess near transfer effects, we divided outcomes in short-term memory (STM) and WM, as the majority of WM definitions recognize both passive storage and active processing as parts of WM (Cowan,

2017). Additionally, correlations within verbal or spatial domains are higher compared to correlations between domains (verbal/visuospatial) (Cowan, 2017; Oberauer et al., 2018). Accordingly, we divided WM outcomes in verbal and visuospatial categories. Reasoning was adopted as a far transfer outcome due to its strong relationship with WM and due to the fact that it is a commonly used measure in the field (Conway et al., 2003; Oberauer et al., 2008). Given that neuropsychological test outcomes varied across studies, they were grouped into broader domains to allow comparisons across studies. A description of each cognitive domain and the corresponding measures is available in app table 6 and app table 7. A minimum of four articles was necessary to compose a category. For verbal WM, the outcomes were grouped into three categories: bwd simple span; complex span; updating. Visuospatial WM had only the bwd simple span category. For STM, only the category “simple span” was created. Reasoning outcomes were grouped according to the tests used to assess reasoning abilities (e.g., Cattell, Raven Standard Progressive Matrix – RSPM, RAPM, Leistungsprüfungssystem Subtest – LPS).

### 2.3. Multilevel-meta-analysis

Effect sizes were calculated to estimate the transfer effect difference between WMT and control condition. The effect sizes of post-intervention and follow-up gains were calculated using Hedges'  $g$  (Hedges, 1989). Since the design used in the individual studies of this meta-analysis have a *pre-posttest control* design, we followed the discussion presented by Morris (2008, p. 369) to calculate the effect sizes measures. More precisely, we used the standardized mean difference described in formula 5, which was originally defined by Becker (1988):

$$g = c(n_E - 1) \frac{M_{post,E} - M_{pre,E}}{SD_{pre,E}} - c(nc - 1) \frac{M_{post,C} - M_{pre,C}}{SD_{pre,C}},$$

where  $M_{pre,E}$  and  $M_{post,E}$  are the experimental group pretest and posttest means,  $SD_{pre,E}$  is the standard deviation of the pretest scores,  $c(m)$  is a bias correction factor,  $n_E$  is the size of the experimental group, and

$M_{pre,C}$ ,  $M_{post,C}$ , and  $nc$  are analogous values for the control group. The bias correction factor is presented in formula 22 as described in Morris (2008, p. 372):

$$c(m) = \sqrt{\frac{2}{m}} \frac{\Gamma(m/2)}{\Gamma((m-1)/2)},$$

where  $\Gamma$  is the gamma function. The sampling variances were obtained through equation 13 of Becker (1988). All effect sizes and sampling variances were automatically computed using the R package “metafor”.

Unfortunately, accurate estimation of the effect size variance in this formula requires the correlation between pre- and posttest scores, which was not available for most of the studies. Therefore, as recommended by Borenstein (2009), a range of plausible correlations ( $r = 0.3, 0.5, 0.7$ ) was considered, and a sensitivity analysis was conducted to ensure that the conclusions from the meta-analysis were robust. A table for the sensitivity analysis is provided in the supplementary material (see table C).

In some studies, more than one measure for the same category was adopted within the same experiment (e.g., Cantarella et al., 2016 reported on two reasoning measures: Cattell and RSPM). In those cases, a multilevel model was adopted for handle multiple effect sizes from the same sample. Using a robust method for dealing with multiple outcomes, such as in the multilevel model or the RVE, is important to avoid bias in the estimates of the effects, standard errors and variances (Moeyaert et al., 2017; Morris, 2008).

Considering that effect sizes from the same study are dependent on

one another, a multivariate meta-analysis is recommended to model these dependencies (Harbord, 2011). Indeed, classic meta-analytic models assume independence among effect sizes. However, this assumption is not realistic with clustered data, such as multiple outcomes from the same study. Multilevel models allow for model dependencies due to clustering and are therefore recommended to account for non-independence in the observed outcomes. Classic meta-analytic models can be considered 2-level models, with participants at level 1 and effect sizes at level 2, whereas multilevel models, also called 3-level models, include clusters at level 3.

In this work, we used multilevel modelling that was complemented with both a sensitivity analysis and the RVE method. Specifically, this procedure consisted of two main steps. First, a full sampling variance-covariance matrix was imputed through the function “impute\_covariance\_matrix” from the “clubSandwich” package by selecting the studies to be the clusters, and the intra-experiments correlation to be 0.5. Second, the corresponding multilevel multivariate random-effects model was assessed through the function “rma.vm” in the “metafor” package. Unstructured correlation matrices were used to allow random effects to be correlated and to have different variances for each outcome. To ensure robustness of the meta-analysis results, complementary analyses were performed through the RVE method and a sensitivity analysis with different correlations ( $r = 0.3, 0.5, 0.7$ ). Robust results have been obtained through the function “coef\_test” from the “clubSandwich” package, following the cluster robust estimator for multivariate/multilevel meta-analytic models described in Hedges et al. (2010). Due to consistent findings observed with these complementary methods, further mixed effects multilevel modelling (using moderators) was only performed for  $r = 0.5$  and  $r = 0.5$  ( $r$  is the pre-posttest correlation and is the intra-experiment correlation).

The significance of the pooled effect size was determined using a  $Z$  test. Effect size was also compared to a  $t$ -test with the Satterthwaite correction (Pustejovsky, 2017) and to a likelihood ratio test based on Skovgaard's statistic (Guolo and Varin, 2012) to confirm the validity of the findings. The effect size for each construct is presented in Table 1. Forest plots with the distribution of effect sizes were then generated for all constructs and categories (see app. Fig. A2). Visual inspection of graphs, Cochran's  $Q$  test, and the  $I^2$  Index were used to assess heterogeneity in random-effects models. The variance components  $\sigma^2_1$  and  $\sigma^2_2$  were used to assess between- and within-studies heterogeneity, respectively, in the multilevel analysis. To address the small number of studies included in some of the analyses, two small sample corrections were performed: Satterthwaite  $p$ -values from the RVE (Pustejovsky, 2017), and Skovgaard's  $p$ -values from second-order likelihood inference (only for 2-level random effects) (Guolo and Varin, 2012).

### 2.4. Influential outcomes

Influential outcomes are considered outliers that exert a strong influence on the results. To ensure the robustness of the results, influential outcomes were removed from each group with at least four outcomes. They were identified by the function “influence” from the “metafor” package, and they are summarized in app. table 8. The analysis of influential studies identified 17 influential outcomes, which were eliminated from the original database.

### 2.5. Moderator analysis

A moderator analysis was conducted with predictors selected from previous literature (Borella et al., 2017a; Bürki et al., 2014; Lilienthal et al., 2013; Stepankova et al., 2014; Verhaeghen et al., 1992; Zinke et al., 2014), considering their influence in visuospatial and verbal WM



Table 1

Effects of working memory training compared with control group by construct.

												Construct	No. of effects (k)	No. of studies (clusters)	RE mean				
															Estimate	95% CI	p-value		
POSTTRAINING	Construct	No. of effects (k)	No. of studies (clusters)	Estimate	RE mean			Q-test	F (%)	$\tau^2$	$\sigma^2_1$	$\sigma^2_2$							
					95% CI	p-value	Skovgaard's p-value	RVE p-value											
	Reasoning	33	24	0.10	[-0.026,0.233]	.12	NA	.13	28.53	11.51	NA	0.01	<0.01	Reasoning	33	24	0.10	[-0.026,0.233]	.12
	Verbal WM	40	20	0.23	[0.065,0.392]	.006 **	NA	.01*	88.79 ***	56.13	NA	<0.01	0.09	Verbal WM	40	20	0.23	[0.065,0.392]	.006 **
	Visuospatial WM	13	10	0.23	[0.029,0.426]	.025 *	NA	.06^	16.03	17.83	NA	0.02	<0.01	Visuospatial WM	13	10	0.23	[0.029, 0.426]	.025 *
FOLLOW-UP	Verbal STM	12	11	0.16	[-0.045,0.363]	.13	NA	.16	12.41	14.07	NA	<0.01	0.01	Verbal STM	12	11	0.16	[-0.045,0.363]	.13
	Visuospatial STM	6	5	-0.03	[-0.388, 0.324]	.86	NA	.74	09.06	45.24	NA	<0.01	0.08	Visuospatial STM	6	5	-0.03	[-0.388, 0.324]	.86
	Reasoning	12	10	0.13	[-0.085, 0.347]	.24	NA	.27	9.36	6.37	NA	0.01	<0.1	Reasoning	12	10	0.13	[-0.085, 0.347]	.24
	Verbal WM	17	9	0.23	[0.006, 0.457]	0.04 *	NA	.08^	18.59	16.35	NA	0.01	0.01	Verbal WM	17	9	0.23	[0.006, 0.457]	.04 *
	Visuospatial WM	11	8	0.14	[-0.089, 0.368]	.23	NA	.14	6.04	<0.01	NA	<0.01	<0.01	Visuospatial WM	11	8	0.14	[-0.089, 0.368]	.23
Verbal STM	6	6	0.18	[-0.097, 0.452]	0.205	0.983	.19	3.85	<0.01	<0.01	NA	NA	Verbal STM	6	6	0.18	[-0.097, 0.452]	.205	
Visuospatial STM	6	5	-0.04	[-0.334, 0.245]	0.763	NA	.72	3.17	NA	NA	<0.01	<0.01	Visuospatial STM	6	5	-0.04	[-0.334, 0.245]	.763	

			RE mean	Q-test	I <sup>2</sup> (%)	$\tau^2$	$\sigma^2_1$	$\sigma^2_2$
			Skovgaard's p-value	RVE p-value				
POSTTRAINING	NA	.13	28.53	11.51	NA	0.01	< 0.01	
	NA	.01*	88.79 ***	56.13	NA	< 0.01	0.09	
	NA	.06^	16.03	17.83	NA	0.02	< 0.01	
	NA	.16	12.41	14.07	NA	< 0.01	0.01	
	NA	.74	09.06	45.24	NA	< 0.01	0.08	
FOLLOW-UP	NA	.27	9.36	6.37	NA	0.01	< .01	
	NA	.08^	18.59	16.35	NA	0.01	0.01	
	NA	.14	6.04	< 0.01	NA	< 0.01	< 0.01	
	0.983	.19	3.85	< 0.01	< 0.01	NA	NA	
	NA	.72	3.17	NA	NA	< 0.01	< 0.01	

Note. ^ $p < .1$ , \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ . NA – Not applicable (only for groups from the same experiment). I<sup>2</sup> – total heterogeneity / total variability;  $\tau^2$  – estimated amount of total heterogeneity;  $\sigma^2_1$  – Variance component of the 3-level model for the between-studies heterogeneity;  $\sigma^2_2$  – Variance component of the 3-level model for the within-studies (effects within studies) heterogeneity. RVE – Robust variance estimation. Number of studies may be smaller than number of effects because each study may have more outcomes for the same construct. RVE and Skovgaard's (only for 2-level random effects) were applied as a sensitivity analysis to check the robustness of the model. P-values did not differ substantially across these analyses indicating the validity of the model.

and STM, as well as, reasoning for both immediately after training and at follow-up. The following variables have been tested as moderators: 1) type of control (active or passive); 2) mean age of the participants; 3) training dose (total number of training in hours); 4) training length (in weeks); 5) total number of sessions; 6) training type (single i.e., complex/simple span, updating training or mixed training, i.e., combination of more than one type of WM task); 7) years of formal education; 8) category of the outcome (e.g., Cattell; RAPM; RSPM); 9) general cognitive ability (measured by the vocabulary test); 10) baseline performance. In this analysis, we used a 3-level random-effects model to assess the overall effect of WMT in post-test and follow-up for each construct, including each moderator separately.

## 2.6. Publication bias

To assess sensitivity to publication bias in this meta-analysis, different complementary methods were utilized, namely, tests for funnel plot asymmetry (Egger et al., 1997), the trim-and-fill method (Duval and Tweedie, 2000a, 2000b), and the Henmi and Copas method (Henmi and Copas, 2010). The sensitivity analysis of the results was investigated with the “leave-one-out method”. Given that publication bias is based on the symmetry of the distribution of the effect sizes in a funnel plot, if we compare very different measures, the distribution is not expected to be symmetric, and it may mislead the analysis. In our sample, studies adopted a large variety of tasks to measure the same construct. Thus, publication bias and the “leave-one-out” method were assessed by the categories of measures. Moreover, asymmetry of funnel plot was verified only in categories having at least 10 experiments, following literature recommendations (Sterne et al., 2011; Zhou et al., 2017). Finally, as these methods aim to identify significant differences between individual experiments, no more than one outcome per experiment can be included in a single plot. Therefore, for groups having at least two outcomes from the same trial, all possible combinations of subgroups, including exactly one outcome per trial, were considered to assess publication bias and the “leave-one-out” method. Funnel plots with the effect sizes of the included studies in all comparisons can be found in the supplementary material section (see Fig. A).

## 3. Results

The results are described in four major sections. First, we describe the different studies that were included in the analysis. Second, we present the small-study effect analyses. The third section targets the main aim of this review which was to verify the WMT effectiveness at posttest and follow-up together with the moderator analysis. Finally, the risk and publication bias results are presented.

### 3.1. Characteristics of included studies

We identified 300 studies (after removal of duplicates), from which 217 were excluded after reading the abstract and 59 after the full-text analysis. Criteria for paper exclusion: a) review paper; b) sample of non-human animals; c) young participants or elderly but not cognitively healthy participants; d) training does not exclusively target WM; e) the active control group performed a WM task; f) absence of control group; g) studies whose sample has been previously used in another study already included in the meta-analysis; h) WMT coupled with tDCS; i) incomplete data. Twenty-four articles (27 experiments) met the inclusion criteria (for a list of the included papers, see table B in the supplementary material) and were selected for the quantitative analysis, which included data for up to 1130 participants. All trials were published in the last ten years, with *Psychology and Aging* as the journal with

the highest number of publications.

The mean age of the participants ranged from 62.9–87.1 years ( $M = 69.5$ ,  $SD = 4.9$ ), and years of formal education ranged from 6 to 17 ( $M = 12.7$  years,  $SD = 2.85$ ). Of these studies, 79% were carried out in Europe ( $n = 19$ ), with the remainder conducted in North America ( $n = 3$ ; 13%) and Asia ( $n = 2$ ; 8%). On average, studies implemented 12 training sessions ( $SD = 8.59$ ; range = 3–40), corresponding to seven total hours ( $SD = 4.36$ ; range = 1.5–20), with a mean session duration of 42 min ( $SD = 13.8$ ; range = 20–60), and an average of three days of sessions per week ( $SD = 1.36$ ; range = 2–7). Follow-up was reported in eight papers, with a mean of eight months after training ( $SD = 4.4$ ; range = 3–18). The completion rate for the whole sample ranged from 70 to 100%. Most of the training was performed in laboratory settings ( $n = 16$ ); however, six trials were conducted at participants' homes. This information was not detailed in three papers (Richmond et al., 2011; Xin et al., 2014). In eight studies, participation was voluntary, one study included both pay and voluntary participation, ten articles reported financial compensation, and five papers did not mention this information.

Regarding the type of trained task (see app. table 4), studies were grouped into three major categories (Schmiedek et al., 2009; Shipstead et al., 2012): complex or simple span task; updating; mixed (i.e., participants were trained on more than one type of WM task). Eight studies included a complex span task, participants were trained on a simple span task in one study (Zinke et al., 2011), and updating training was observed in ten studies. Five studies had mixed training. Regarding the modality of training (verbal vs. visuospatial), 10 studies included training with verbal stimuli, five included training with visuospatial stimuli, and the remaining nine were crossmodal. All studies, except Pergher et al. (2018); Xin et al. (2014); Zajac-Lamparska and Trempala (2016), had adaptive training. Fourteen articles had an active control group, while ten had a passive control group (PCG). As seen in app. table 3, characteristics regarding type of training and control, outcomes and follow-up varied across studies.

Heterogeneity indexes among studies in the different analyses were low to moderate (Higgins et al., 2003). However, we opted for the random model considering the clinical and methodological heterogeneity found among studies (Higgins and Green, 2008). Before proceeding to the meta-analysis, small-studies effects were explored. The comparison between random-effect modelling, fixed-effect modelling and the Henmi and Copas method were conducted to address this issue. The results of this analysis are summarized in app. table 9. The conclusions of the three models produced very similar results, and in 71% of the cases the difference was  $\leq 0.001$ , not affecting the significance of the results. The most distinct case happened for verbal complex span at posttest, for which the mean effect from the random-effects model was 0.34, 95% CI = [0.09, 0.58], and the common effect from the fixed-effects model was 0.31, 95% CI = [0.14, 0.49]. In both cases, CI did not include zero, confirming its statistical significance. Additionally, sensitivity analysis confirmed that the meta-analytic findings were robust regarding the tested correlation coefficients. Indeed, by visual inspection of the table C in the supplementary material, it is possible to observe that when the correlation is assumed to be lower, at  $r = .3$ , or higher, at  $r = .7$ , the estimated summary effect varies by no more than 0.04.

### 3.2. WMT efficacy and moderator analysis

In this section the results from the effect of WMT on transfer task immediately after training (aim 1) and at follow-up (aim 2), as well as, a moderator analysis (aim 3) will be presented. Results from the classical  $p$ -value or those corrected for small samples (Skovgaard's and

RVE) did not differ considerably, so we reported the multi-level  $p$ -value in the text and all the values in Table 1. The comparisons only had a small difference between the multi-level  $p$ -value ( $p = .03$ ) and the RVE ( $p = .06$ ) for visuospatial WM in posttest and the multi-level  $p$ -value ( $p = .04$ ) and RVE  $p$ -value ( $p = .08$ ) for verbal WM at follow-up. Therefore, the results regarding visuospatial WM in posttest and verbal WM at follow-up should be interpreted with more caution.

We did not find any significant difference between the control types (passive versus active control groups) in the moderation analysis (see app table 10), except for visuospatial WM at posttest. Additionally, we performed a sensitivity analysis, running the analysis separately for passive and active control groups. The comparison with both passive and active control group merged did not yield an effect size greater than when we performed the comparison of experimental group with studies that included only an active control group, except for visuospatial WM at posttest. Many of the included trials had passive control group ( $n = 10$ ). If we had excluded those trials from the analyses, some of the comparisons would have a very few studies, decreasing the power of the analyses. Accordingly, the results from both control groups were merged into a single control condition. The effect sizes were calculated comparing the experimental condition with the merged control condition.

### 3.2.1. Aim 1: examining the generalization of training effects to non-trained tasks (near and far transfer)

WMT effects were examined on near transfer constructs (visuospatial and verbal WM, and visuospatial and verbal STM) as well as on a far transfer construct (reasoning) immediately post-training.

Verbal WM: A significant transfer effect was identified for verbal WM (0.23; 95% CI [0.07, 0.39]).

Visuospatial WM: A significant transfer effect was identified for WM in the visuospatial modality (0.23; 95% CI [0.03, 0.43]).

Verbal and visuospatial STM: No significant transfer effects were identified for verbal (0.16; 95% CI [-0.05, 0.36]) or visuospatial STM (-0.03; 95% CI [-0.39, 0.32]).

Reasoning: For reasoning, the effects were not significant ( $p = .08$ ) at posttest (0.10; 95% CI [-0.03, 0.23]).

### 3.2.2. Aim 2: verifying the maintenance of the effects at follow-up

Concerning the long-term effects of WMT, we observed that the effects were also observed during follow-up to verbal WM (0.23; 95% CI [0.01, 0.46]). However, in visuospatial WM analysis, the effect was not

significant (0.14; 95% CI [-0.09, 0.37]). Regarding reasoning, results were also not significant (0.13; 95% CI [-0.09, 0.35]), as well as for verbal STM (0.18; 95% CI [-0.10, 0.45]) and visuospatial STM (-0.04; 95% CI [-0.33, 0.25]).

### 3.2.3. Aim 3: testing moderator variables

Here we examined if the variable age, training dose, number of sessions, training type, training duration, years of formal education, vocabulary score, baseline performance and type of outcome might moderate training effects. The results are presented in Table 2. The moderator analysis was significant ( $p < .05$ ) for number of sessions, training length (in weeks) and training dose (in hours), i.e., the gains in reasoning and verbal STM immediately after training are small when training duration increases. Additionally, while the effect of WMT on verbal STM was linearly moderated by training hours and training length, the effect of WMT on Reasoning-posttest was also moderated by the former factors together with the number of sessions. Table 2 outlines these moderator roles. Indeed, the approximation by higher polynomial degrees were also assessed but, in each case, no advantage over the linear approach was observed. Specifically, no asymptotic behaviour was detected, as such characteristic would imply a significant variation in the rate of change of the WMT effect with respect to the corresponding independent variable.

Regarding the training type, we observed that the studies that included mixed training (i.e., having more than one type of WM tasks) had smaller effects on reasoning immediately after training than the training of updating or complex span tasks alone. Additionally, studies having the Cattell Test as an outcome displayed a higher gain than studies that used other measures at posttest (RAPM; RSPM; LPS). For verbal WM, the gains were higher in complex span tasks than in simple span and updating tasks at posttest. Type of control group was a significant moderator for verbal WM at posttest, with the effect size of studies using a passive control group being higher than studies that used an active control group. Finally, baseline performance moderated the effects on visuospatial STM at immediate posttest, with participants with lower performance showing more benefits with the training.

In summary, WMT had a small significant and long-lasting effect on verbal WM (specifically on complex span outcomes). For visuospatial WM, gains were only observed at posttest, but not at follow-up. Far transfer for reasoning was not observed. Training length, number of sessions, training dose (total training duration in hours), type of training and adopted outcomes (Cattell; and complex span), type of

**Table 2**  
Moderator effects (significant results).

Construct	Moderator effect	Estimate	SE	p-value	QE	QM – Test of moderators	$\sigma^2_1$	$\sigma^2_2$
Reasoning at immediate posttest	Measure – Cattell	0.39	0.14	.005**	20.70	7.82 **	< 0.01	< 0.01
	Training dose (hours)	-0.04	0.01	.001 **	17.46	11.040 ***	< 0.01	< 0.01
	Number of sessions	-0.02	0.01	.004**	20.17	8.35**	< 0.01	< 0.01
	Training length (in weeks)	-0.11	0.04	.004 **	20.38	8.15 **	< 0.01	< 0.01
	Training Type – Mixed	-0.41	0.13	.001**	18.36	10.16 **	< 0.01	< 0.01
Verbal WM at immediate posttest	Measure – Complex span	0.27	0.13	.046 ***	80.67	4.00 *	< 0.01	0.08
Visuospatial WM at immediate posttest	Control – PC – AC	0.54	0.24	.023*	10.86	5.17 *	< 0.01	< 0.01
Verbal STM at immediate posttest	Training dose (in hours)	-0.04	0.02	.043*	8.33	4.08*	< 0.01	< 0.01
	Training length (in weeks)	-0.11	0.05	.033*	7.89	4.53*	< 0.01	< 0.01
Visuospatial STM at immediate posttest	Baseline performance	-0.06	0.02	.01*	2.33	6.73**	< 0.01	< 0.01

Note. \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ ;  $\sigma^2_1$  – Variance component of the 3-level model for the between-studies heterogeneity;  $\sigma^2_2$  – Variance component of the 3-level model for the within-studies heterogeneity. QE – test for residual heterogeneity when moderators are included. QM – test statistic for the omnibus test of coefficients. Moderator effects with non-significant results were not presented, they were mean age of the participants, years of formal education, vocabulary performance. Analyses of follow-up did not have any significant moderator.

control group and baseline performance appeared as significant moderator variables at posttest assessment.

### 3.3. Publication and risk of bias

Assessment of risk of bias is important when performing a review because it is an index of the quality of included data, and it could also explain heterogeneity when it is highly observed (Viswanathan et al., 2008). Two authors independently assessed the risk of bias. In general, we observed a substantial absence of information for most studies, which limited the ability to classify the risk of bias. Considering the randomization processes (selection bias), 19% of the studies presented risk of bias, whereas in 74% the risk of bias was not clear. Seven percent of the studies adequately reported random sequence generation. Regarding allocation concealment, 22% presented a high risk of bias, 7% adequately reported data, and the remaining 70% did not report on allocation concealment. For blinding (performance bias), 30% of the studies had low risk of bias (compared with 30% with high risk), and 40% of the studies did not mention blindness procedures. Seventy percent of the studies did not exclude data from participants who dropped out or with missing data. Fifteen percent had high risk of incomplete outcome data, while this was not clear in 15% of the studies. Generally, the studies had high completion rates (ranging from 86% to 100%), although the completion rate was not clear for all studies. Similarly, most articles (93%) reported all outcomes, although they did not state which outcome was the primary. Seven percent presented high risk of selective reporting. Additionally, the lack of adequate correction for multiple comparisons and for baseline group differences were other potential bias observed here. Another possible source of bias was the lack of appropriate screening measures of cognitive decline and of affect disorders such as anxiety and depression. A summary graph of the risk of bias is displayed in app. Fig. 3.

Analysis of publication bias assesses if the set of evidence is biased due to the fact that positive findings are more likely to be published. The analysis of several methods of publication bias (trim-and-fill, leave-one-out, asymmetric tests, and Hemni and Copas) suggested a small presence of publication bias, although it did not seem to substantially alter the results. Trim-and-fill is a method that estimates the number of studies missing in the funnel plot (Duval and Tweedie, 2000b). It was only used in analyses with at least 10 studies; otherwise, the test would not have sufficient power to verify asymmetry (Sterne et al., 2008; Zhou et al., 2017). This analysis suggested the presence of publication bias in only two cases (simple span and complex span at posttest). Additionally, given that the big issue of publication bias is that the positive results are more representative in the published literature (Mlinarić et al., 2017), it is important to highlight that trim-and-fill method identified only two cases of missing studies (verbal simple span STM and verbal updating WM, both at posttest), however the effect sizes of the corresponding categories were not significant in verbal simple span STM and verbal updating WM at posttest.

The leave-one-out method was performed by a sensitivity analysis where one study at a time was removed from the analysis to verify the influence of a single study in the finding. This method showed sensitivity of results to individual studies in three cases (verbal fwd simple span at posttest; Cattell and verbal complex span at follow-up). However, in the first two cases, the elimination of a unique experiment would cause a significant pooled effect size, while only for complex span the elimination of a study (among three) would cause a non-significant result. Asymmetric tests indicated publication bias in only one case (verbal simple span at posttest), the same comparison already

identified with the trim-and-fill method. Finally, in all cases, the Hemni and Copas robust estimation was not significantly different from the random-effects results, showing that publication bias did not change the overall meta-analytic effects in a significant manner. Therefore, the positive effect of publication bias was not a big issue here.

Overall, the presence of bias did not seem to influence the results as supported by the former publication bias methods (see app. table 9), as well as, by the similarity between effect sizes of studies that presented more criteria classified as high risk of bias (see app. Fig. 3) (e.g., Goghari and Lawlor-Savage, 2017; Heinzel et al., 2016; Stepankova et al., 2014; Zinke et al., 2011) and those having a lower risk of bias (e.g., Borella et al., 2017b, 2013; Guye and von Bastian, 2017; Lange and Süß, 2015; Weicker et al., 2018).

## 4. Discussion

This meta-analytical review aimed to verify the gains of WMT on transfer measures in healthy older adults. In contrast to previous meta-analyses, we used different analytical methods to address multiple outcomes and the lack of correlation reports. Additionally, a description of the studies included in the review is provided along with a comprehensive overview of different studies in the WMT field.

The high variability between the experiments challenged data aggregation and, consequently, data interpretation. The studies presented different experimental and control tasks (see app. table 4 and app table 5), different outcomes (see app table 7), and training protocols. Follow-up also varied broadly across trials, although it was seldom included in the experimental protocol (see app table 3).

Regarding the results of the effectiveness of WMT at posttest (aim 1), participants assigned to a WMT group displayed a small significant near transfer effect size of 0.2 for verbal and visuospatial WM, compared to the participants who received a placebo or non-intervention. These results are in line with previous meta-analyses that have shown small to medium near effect sizes immediately after training (Karch and Verhaeghen, 2014; Melby-Lervåg et al., 2016; Melby-Lervåg and Hulme, 2013). For example, Karch and Verhaeghen (2014) observed a small near effect size of 0.3 after removal of publication bias (trim-and-fill method). We also observed that WMT had no significant impact on STM, which conflicts with the results of previous research (Schwaighofer et al., 2015). These differences among studies may be due to methodological differences, as Schwaighofer and colleagues (2015) included older adults as well as children and young adults. Moreover, it might be the case that the lack of effect in STM may be due to a preservation of this ability with age (Nittrouer et al., 2016; Olson et al., 2004). Therefore, there is less room for transfer in this ability after WMT. Nevertheless, this hypothesis needs to be further explored as there was one study showing a strong positive effect of WMT on STM (Heinzel et al., 2013). As we observed in the moderator analysis, variables such as the training dose and length, as well as, baseline performance interfered with the effects, which may cause heterogeneity across studies. For the reasoning, there was no significant transfer effect. In fact, a previous meta-analysis (Karch and Verhaeghen, 2014) only yielded a "marginally significant" far transfer effect that was not fully corroborated by our study with a greater number of WMT trials included in the analysis.

With respect to the WMT long-term effects (aim 2), only ten studies reported follow-up assessments; therefore, the results should be considered with caution. Near transfer effects seem to be maintained at follow-up only for verbal WM. These results are in agreement with Schwaighofer et al. (2015) and partially consistent with Melby-Lervåg



et al. (2016; 2013), who only observed a significant maintenance effect in WM outcomes.

We performed a moderator analysis with the following variables as moderators of transfer effects on STM, WM and reasoning at posttest and follow-up (aim 3): 1) type of control (active/passive); 2) the mean age of participants; 3) training dose (total number of training in hours); 4) training length (in weeks); 5) number of training sessions; 6) training type (single: complex span or updating; mixed training: more than one type of WM task); 7) years of formal education; and 8) category of the outcome (e.g., Cattell; RAPM; complex span); 9) vocabulary score; 10) baseline performance. The variables that explained heterogeneity of the effect sizes in reasoning at posttest were the category of the outcome (i.e., Cattell), training length/dose, number of training sessions, and training type (i.e., mixed training). For verbal WM at posttest, the category of the outcome (i.e., complex span) was the variable that explained heterogeneity of the effect sizes. This means that studies having complex span as outcome found more positive effects than studies using another WM measures. For visuospatial WM at posttest, the type of control group (active versus passive) was a significant moderator, with studies using passive control groups presenting higher effect sizes. For verbal STM at posttest, training length and hours were the significant moderators. For visuospatial STM at posttest, baseline performance moderated the results, with participants with lower performance gaining more with the training.

The fact that some measures (i.e., Cattell Test and Complex Span Task) displayed more significant effect sizes than others in the moderator analysis highlights the role of the measures to evaluate the training effects. For reasoning, the effect size on the Cattell Test was significant, showing a positive moderation effect of this test on far transfer. This result is in line with the results of previous reviews which showed a slightly larger effect of the Cattell Test compared to Raven's Test (Mansur-Alves and Silva, 2017). This finding could be explained by the fact that the Cattell Test consists of different subtests (series, analogies, matrices and classification), which may position it as a more complete indicator of reasoning compared to tests that only have figural type items (e.g., Raven), as postulated by Gignac (2015). Furthermore, this result is consistent with the claim of Shipstead et al. (2012) re-garding the importance of having different instruments to assess transfer effects in the experiments, ensuring that all facets of the construct are assessed.

Considering the moderation effect of training dose/length, either in reasoning or verbal STM, we found unexpected results. For both variables, the results showed a significant negative effect, i.e., that more training duration (total number of hours and length) produced smaller effect sizes. Other variables probably influenced this analysis, such as the type of training performed: most of the shorter duration studies applied the same training task which may be more effective than the training adopted by the long-duration studies (Borella et al., 2017a). It is also noteworthy that only one study had higher dosages of training (more than 15 h) (Goghari and Lawlor-Savage, 2017), whereas six out of twenty had only three sessions (Borella et al., 2017b, 2014, 2013, 2010; Cantarella et al., 2017b, 2017a). Previously, Karbach and Verhaeghen (2014) and Melby-Lervåg and Hulme (2013) failed to find a significant influence of total training duration in effect size, except for one measure, the Stroop task in Melby-Lervåg and Hulme (2013). In contrast, Schwaighofer et al. (2015) found a positive influence of total training duration on visuospatial STM and of session duration on verbal STM. Weicker et al. (2016) documented a positive correlation between the number of sessions and the effect sizes. In this case, the authors compared two groups (> 20 sessions vs. < 20 sessions) and observed

that more training sessions produced larger effect sizes. Nonetheless, the total number of hours was not related to the effect size (> 10 h vs. < 10 h). Finally, similar to our results, a previous meta-analysis on video-game training have shown that short training produced stronger effects than long training (Toril et al., 2014). These discrepant findings need to be further addressed in new randomized controlled trials.

Other factors such as motivation and performance anxiety should also be considered (Delphin-Combe et al., 2016; Jaeggi et al., 2014). As participants are older adults, some of them may be unfamiliar with the use of computers (most of the trainings are computerized), and long training durations may lead to demotivation (Laguna and Babcock, 1997). Additionally, participants might not be receptive to extensive training because the training would compete with their other activities for time. Another finding worth considering is the fact that mixed training negatively moderated the effects on reasoning. In other words, the experience of different tasks in the same programme may be less effective than repeating the same task or similar tasks during the training (for similar results, see von Bastian et al., 2013). Perhaps targeting a specific process during training yields sizeable gains, whereas the training of multi-WM processes may lead to a competition for re-sources that underpin the transfer effects.

In short, considering the aim 1 (effectiveness at posttest), our results supported only the presence of near transfer effects. For the aim 2 (effectiveness at follow-up), our results supported the maintenance of near transfer effects only on verbal WM. For the aim 3, our data suggested that the type of outcome (Cattell and complex span), total training duration/length/number of sessions, baseline performance, type of control group and type of trained task (mixed task) moderate the transfer effects.

Melby-Lervåg and Hulme (2016) identified two main problems with previous meta-analyses that showed promising effects of WMT (Au et al., 2015; Karbach and Verhaeghen, 2014). The first was related to the calculation of a mean effect size without considering the baseline performance. It is noteworthy, however, the absence of correlations between baseline and posttest assessment in the original papers challenges the calculation of the Hedge's *g* change variance. To address this issue, in this review, the effect size calculation was based on the pre- to posttest score difference (Borenstein et al., 2009; Morris, 2008), and we also ran a sensitivity analysis with different values of correlation coefficients.

The second problem pointed out by the authors was the importance of comparing studies with active versus passive control groups. To address this, we performed a moderator analysis with the type of control as moderator which showed a significant effect only for visuospatial WM at posttest (see App Table 10). We also ran a sensitive analysis with active and passive control group separately (see supplementary material, table D). The effect sizes did not change considerably from the previous results with the merged control group. The exception was the visual WM at posttest in which the results became insignificant. In this analysis, results from RVE and multi-level *p*-value also differed from each other showing that this finding needs further evidence. Moreover, it is noteworthy that one influential study with a big positive effect size (Borella et al., 2014 – experiment 1) was excluded. If we had kept this study, the analysis would be significant either case. Probably there is in fact an effect in the visuospatial WM, however given the inconsistency in different analysis, it is not possible to draw a clear conclusion.

In contrast, in the study of Melby-Lervåg and Hulme (2016), some of the meta-analytical results changed when the analysis was performed separately for active and passive control groups. Our findings partially corroborated the results of Weicker et al. (2016), Melby-Lervåg and

Hulme (2013), and Karbach and Verhaeghen (2014) that did not find a significant influence of the type of control condition in the outcomes. It is noteworthy, however, that Melby-Lervåg and Hulme (2016) had a diversified sample, including a broader range of ages and learner status within the same analysis, which may explain the differences found.

Relatively to the assessment of risk of bias, most authors did not report data regarding random sequence generation, allocation concealment, and blinding. Among the other risks of bias identified, some trials have performed multiple outcome comparisons without correction or did not use validated screening measures of cognition and affect. Other experiments showed differences between conditions at baseline, most likely due to inappropriate randomization. Some studies were exploratory, not stating primary/secondary analysis, nor including a priori sample size calculations. Nonetheless, in the current review, the risk of bias was not problematic since the same pattern of results was found both in studies that fulfilled most of the criteria and in studies that satisfied only a few. Additionally, more recent studies considered this limitation and implemented a more appropriate experimental design (Guye and von Bastian, 2017; Weicker et al., 2018).


The primary limitation of this review is the fact that we pooled different methodological studies together. However, we have done moderator analyses and combined outcomes in categories to address this variability. Second, although we considered a Ph.D. thesis, we did not perform an extensive grey literature search, which may have introduced publication bias in our analysis. It is noteworthy, however, that publication bias analysis did not indicate a strong presence of such bias, especially regarding positive statistical effects. Third, in some comparisons, we had a low number of trials included ( $n < 10$ ), especially with follow-up analyses. Fourth, some of the included studies had a small sample in each comparison ( $n < 20$ ). Even though, this limitation was addressed in the analysis by applying corrections for small samples to the effect size calculation. Fifth, two studies were not included due to the lack of replies from the contacted authors (missing data). Finally, our results may not be valid for the whole ageing population because most studies were conducted with a selective population. To illustrate, most trials had participants with a high level of schooling ( $M = 12.67$  years), and most of trials had younger older adults as participants ( $M = 69.55$ ). Therefore, additional studies with older populations and participants with lower levels of schooling are needed (e.g., da Silva and Yassuda, 2009; Golino and Flores-Mendoza, 2016).

Finally, some recommendations are suggested for future studies in the WMT field. New trials should address different training formats that

are best suited for the elderly (i.e., optimal session duration, total intervention time and intervals between sessions) (e.g., Penner et al., 2012). Another critical point is related to the importance of increasing the training level of difficulty. In our sample, 95% of the trials were adaptive, meaning that the trained task was adjusted in difficulty according to the participants' performance. However, von Bastian and Eschen (2016) found that participants did not perform better with adaptive tasks than with tasks of self-selected difficulty. Furthermore, a next step could be to compare different WMT programmes as illustrated by Basak and O'Connell (2016), who showed a superiority effect of an unpredictable memory updating training over a predictable one. We also encourage comparisons between web-based interventions and more traditional laboratory approaches (Schwaighofer et al., 2015). Subsequently, researchers should verify how to keep participants engaged in the training programmes. For example, group cognitive trainings could be more motivating than individualized trainings (Kelly et al., 2014). Other approaches such as combining techniques (e.g., non-invasive electrical brain stimulation or physical exercise) could boost WMT effects (Oswald et al., 2006; Teixeira-Santos et al., 2015).

Protocols should be designed to follow participants over more extended periods of time. The outcomes selection could also be rethought. Namely, we could have different outcomes to assess different facets of the same construct (Weicker et al., 2018), and we could account for more clinical relevance and external validity. For example, some promise has been seen regarding the generalizability of results for real life: Cantarella et al. (2016) used everyday problem solving and timed basic daily activities as outcomes; Lange and Süß (2015) had questionnaires for cognitive failures in everyday life; Takeuchi et al. (2014) assessed the effect of WM training on emotional states; and Borella et al. (2019) assessed transfer for everyday life in old-old participants. Eventually, subjective cognitive functioning could be included. Similarly, surrogate outcomes, such as magnetic resonance imaging and electro-physiological recordings, could be used to support the efficacy of the intervention and to define the best training protocol regarding brain plasticity (Buschkuhl et al., 2012; Dahlin et al., 2008b; Heinzel et al., 2016; Takeuchi et al., 2014, 2013).

Other factors that may moderate gains (e.g., motivation; personality; financial compensation) should be further scrutinized (Au et al., 2015; Borella et al., 2013; Zinke et al., 2011). Regarding the population, studies with different age and formal education subgroups are warranted. To the best of our knowledge, no former study in the field has been conducted with illiterate people, mainly because few studies are carried out in developing countries. However, this group is more



**Fig. 1.** Schematic representation of the main findings of the current meta-analysis. X = non-significant results; Solid line = significant results; ↓ = negative moderating effect; ↑ = positive moderating effect.

vulnerable to cognitive decline (Brucki, 2010) and in need of cognitive care opportunities. Finally, regarding risk of bias, future studies should be careful about the proper implementation of the randomization process, allocation concealment, blinding, incomplete outcome and data reporting.

5. Conclusion

Neuroplasticity, the brain and behavioural capacity of restructura-tion according to environmental demands, is verified even in late stages of development (Landi and Rossini, 2010), and WMT has been studied as a promising tool to promote it. Our analysis suggested the general-ization of WMT to near transfer tasks. Far-transfer effects were not verified, except for the studies whose Cattell Test was used to assess reasoning. Moderator analysis did not show the influence of type of control group (active versus passive), except for one comparison: vi-suospatial WM at posttest. Importantly, the adopted measures, type of training, training length and duration, baseline performance were sig-nificant variables moderating the effects sizes. Overall, the general-ization of WMT seems to be limited to the WM construct (see Fig. 1).

Appendix A




Fig. A1. PRISMA flow diagram.




Fig. A2. Posttest Forest plots. Note. Pooled effect size for subcategories (grey diamond) and constructs (blue diamond).

## Appendix B. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.neubiorev.2019.05.009>.

## References

- Au, J., Sheehan, E., Tsai, N., Duncan, G.J., Buschkuhl, M., Jaeggi, S.M., 2015. Improving fluid intelligence with training on working memory: a meta-analysis. *Psychon. Bull. Rev.* 22, 366–377. <https://doi.org/10.3758/s13423-014-0699-x>.
- Bäckman, L., Lindenberger, U., Li, S.-C., Nyberg, L., 2010. Linking cognitive aging to alterations in dopamine neurotransmitter functioning: recent data and future avenues. *Neurosci. Biobehav. Rev.* <https://doi.org/10.1016/j.neubiorev.2009.12.008>.
- Bäckman, L., Waris, O., Johansson, J., Andersson, M., Rinne, J.O., Alakurtti, K., Soveri,

- A., Laine, M., Nyberg, L., 2017. Increased dopamine release after working-memory updating training: neurochemical correlates of transfer. *Sci. Rep.* 7. <https://doi.org/10.1038/s41598-017-07577-y>.
- Ball, K., Berch, D.B., Helmers, K.F., Jobe, J.B., Leveck, M.D., Marsiske, M., Morris, J.N., Rebok, G.W., Smith, D.M., Tennstedt, S.L., Unverzagt, F.W., Willis, S.L., 2002. Effects of cognitive training interventions with older adults: a randomized controlled trial. *JAMA* 288, 2271–2281. <https://doi.org/10.1001/jama.288.18.2271>.
- Basak, C., O'Connell, M.A., 2016. To switch or not to switch: role of cognitive control in working memory training in older adults. *Front. Psychol.* 7, 1–18. <https://doi.org/10.3389/fpsyg.2016.00230>.



- Beatty, E.L., Vartanian, O., Mackey, A.P., 2015. The prospects of working memory training for improving deductive reasoning. *Front. Hum. Neurosci.* 9, 1–2. <https://doi.org/10.3389/fnhum.2015.00056>.
- Becker, B.J., 1988. Synthesizing standardized mean??? Change measures. *Br. J. Math. Stat. Psychol.* <https://doi.org/10.1111/j.2044-8317.1988.tb00901.x>.
- Bisiacchi, P.S., Tarantino, V., Ciccola, A., 2008. Aging and prospective memory: the role of working memory and monitoring processes. *Aging Clin. Exp. Res.* <https://doi.org/10.1007/BF03324886>.
- Borella, E., Carretti, B., Riboldi, F., De Beni, R., 2010. Working memory training in older adults: evidence of transfer and maintenance effects. *Psychol. Aging* 25, 767–778. <https://doi.org/10.1037/a0020683>.
- Borella, E., Carretti, B., Zanon, G., Zavagnin, M., De Beni, R., 2013. Working memory training in old age: an examination of transfer and maintenance effects. *Arch. Clin. Neuropsychol.* 28, 331–347. <https://doi.org/10.1093/arclin/act020>.
- Borella, E., Carretti, B., Cantarella, A., Riboldi, F., Zavagnin, M., De Beni, R., 2014. Benefits of training visuospatial working memory in young-old and old-old. *Dev. Psychol.* 50, 714–727. <https://doi.org/10.1037/a0034293>.
- Borella, E., Carbone, E., Pastore, M., De Beni, R., Carretti, B., 2017a. Working memory training for healthy older adults: the role of individual characteristics in explaining short- and long-term gains. *Front. Hum. Neurosci.* <https://doi.org/10.3389/fnhum.2017.00099>.
- Borella, E., Carretti, B., Sciore, R., Capotosto, E., Taconnat, L., Cornoldi, C., De Beni, R., 2017b. Training working memory in older adults: is there an advantage of using strategies? *Psychol. Aging* 32, 178–191. <https://doi.org/10.1037/pag0000155>.
- Borella, E., Cantarella, A., Carretti, B., De Lucia, A., De Beni, R., 2019. Improving everyday functioning in the old-old with a working memory training. *Am. J. Geriatr. Psychiatry.* <https://doi.org/10.1016/j.jagp.2019.01.210>.
- Borenstein, M., 2009. Effect sizes for continuous data. *The Handbook of Research Synthesis and Meta-Analysis*. pp. 221–235.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T., Rothstein, H.R., 2009. Effect Sizes Based on Means. *Intro to Meta-Analysis*. pp. 21–32. <https://doi.org/10.1002/9780470743386.ch6>.
- Braver, T.S., West, R., 2008. Working Memory, Executive Control, and Aging. *the Handbook of Aging and Cognition*, 3rd ed. .
- Braver, T.S., Cohen, J.D., Nystrom, L.E., Jonides, J., Smith, E.E., Noll, D.C., 1997. A parametric study of prefrontal cortex involvement in human working memory. *Neuroimage.* <https://doi.org/10.1006/nimg.1996.0247>.
- Brehmer, Y., Rieckmann, A., Bellander, M., Westerberg, H., Fischer, H., Bäckman, L., 2011. Neural correlates of training-related working-memory gains in old age. *Neuroimage* 58, 1110–1120. <https://doi.org/10.1016/j.neuroimage.2011.06.079>.
- Brucki, S.M.D., 2010. Illiteracy and dementia. *Dement. e Neuropsychol.* 4, 153–157. <https://doi.org/10.1590/S1980-57642010DN40300002>.
- Bürki, C.N., Ludwig, C., Chicherio, C., de Ribaupierre, A., 2014. Individual differences in cognitive plasticity: an investigation of training curves in younger and older adults. *Psychol. Res.* 78, 821–835. <https://doi.org/10.1007/s00426-014-0559-3>.
- Buschkuhl, M., Jaeggi, S.M., Jonides, J., 2012. Neuronal effects following working memory training. *Dev. Cogn. Neurosci.* <https://doi.org/10.1016/j.dcn.2011.10.001>.
- Cantarella, A., Borella, E., Carretti, B., Kliegel, M., De Beni, R., 2017a. Benefits in tasks related to everyday life competences after a working memory training in older adults. *Int. J. Geriatr. Psychiatry.* <https://doi.org/10.1002/gps.4448>.
- Cantarella, A., Borella, E., Carretti, B., Kliegel, M., Mammarella, N., Fairfield, B., De Beni, R., 2017b. The influence of training task stimuli on transfer effects of working memory training in aging. *Arch. Psychol. Relig. / Arch. Fä/4r Relig.* <https://doi.org/10.1016/j.psfr.2017.04.005>.
- Chan, J.S.Y., Wu, Q., Liang, D., Yan, J.H., 2015. Visuospatial working memory training facilitates visually-aided explicit sequence learning. *Acta Psychol. (Amst).* <https://doi.org/10.1016/j.actpsy.2015.09.008>.
- Clare, L., Woods, R.T., 2004. Cognitive training and cognitive rehabilitation for people with early-stage Alzheimer's disease: a review. *Neuropsychol. Rehabil.* <https://doi.org/10.1080/09602010443000074>.
- Constantinidis, C., Klingberg, T., 2016. The neuroscience of working memory capacity and training. *Nat. Rev. Neurosci.* <https://doi.org/10.1038/nrn.2016.43>.
- Conway, A.R.A., Kane, M.J., Engle, R.W., 2003. Working memory capacity and its relation to general intelligence. *Trends Cogn. Sci.* 7, 547–552. <https://doi.org/10.1016/j.tics.2003.10.005>.
- Cowan, N., 2017. The many faces of working memory and short-term storage. *Psychon. Bull. Rev.* <https://doi.org/10.3758/s13423-016-1191-6>.
- da Silva, H.S., Yassuda, M.S., 2009. Memory training for older adults with low education: mental images versus categorization. *Educ. Gerontol.* 35, 890–905. <https://doi.org/10.1080/03601270902782487>.
- Dahlin, E., Nyberg, L., Bäckman, L., Neely, A.S., 2008a. Plasticity of executive functioning in young and older adults: immediate training gains, transfer, and long-term maintenance. *Psychol. Aging* 23, 720–730. <https://doi.org/10.1037/a0014296>.
- Dahlin, E., Stigsdotter Neely, A., Bäckman, L., Larsson, A., 2008b. Transfer of learning after updating training mediated by the striatum. *Science (80- )* 320, 1510–1512. <https://doi.org/10.1126/science.1155466>.
- Delphin-Combe, F., Bathsavanis, A., Rouch, I., Liles, T., Vannier-Nitenberg, C., Fantino, B., Dauphinaut, V., Krolak-Salmon, P., 2016. Relationship between anxiety and cognitive performance in an elderly population with a cognitive complaint. *Eur. J. Neurol.* 23, 1210–1217. <https://doi.org/10.1111/ene.13004>.
- Diamond, B.J., Deluca, J., Rosenthal, D., Vlad, R., Davis, K., Lucas, G., Noskin, O., Richards, J.A., 1999. Information processing in older versus younger adults: accuracy versus speed. *Int. J. Rehabil. Heal.* <https://doi.org/10.1023/A:1012911203468>.
- Duval, S., Tweedie, R., 2000a. A nonparametric “Trim and fill” method of accounting for publication Bias in meta-analysis. *J. Am. Stat. Assoc.* 95, 89–98. <https://doi.org/10.1080/01621459.2000.10473905>.
- Duval, S., Tweedie, R., 2000b. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication Bias in meta-analysis. *Biometrics* 56, 455–463. <https://doi.org/10.1111/j.0006-341X.2000.00455.x>.
- Egger, M., Davey Smith, G., Schneider, M., Minder, C., 1997. Bias in meta-analysis de-tected by a simple, graphical test. *BMJ.*
- Gignac, G.E., 2015. Raven's is not a pure measure of general intelligence: implications for g factor theory and the brief measurement of g. *Intelligence* 52, 71–79. <https://doi.org/10.1016/j.intell.2015.07.006>.
- Goghari, V.M., Lawlor-Savage, L., 2017. Comparison of cognitive change after working memory training and logic and planning training in healthy older adults. *Front. Aging Neurosci.* 9. <https://doi.org/10.3389/fnagi.2017.00039>.
- Golino, M.T.S., Flores-Mendoza, C.E., 2016. Development of a cognitive training program for the elderly. *Rev. Bras. Geriatr. e Gerontol.* 19, 769–785. <https://doi.org/10.1590/1809-98232016019.150144>.
- Gordon, M., Lumley, T., 2016. forestplot: Advanced Forest Plot Using “grid” Graphics. Guolo, A., Varin, C., 2012. The R package ‘pkmetaLik for likelihood inference in meta-analysis. *J. Stat. Softw.* 50, 1–19. <https://doi.org/10.18637/jss.v050.i07>.
- Guye, S., von Bastian, C.C., 2017. Working memory training in older adults: bayesian evidence supporting the absence of transfer. *Psychol. Aging.* <https://doi.org/10.1037/pag0000206>.
- Harbord, R.M., 2011. Commentary on “Multivariate meta-analysis: potential and promise.”. *Stat. Med.* <https://doi.org/10.1002/sim.4278>.
- Hedges, L.V., 1989. An unbiased correction for sampling error in validity generalization studies. *J. Appl. Psychol.* 74, 469–477. <https://doi.org/10.1037/0021-9010.74.3.469>.
- Hedges, L.V., Tipton, E., Johnson, M.C., 2010. Robust variance estimation in meta-regression with dependent effect size estimates. *Res. Synth. Methods* 1, 39–65. <https://doi.org/10.1002/jrsm.5>.
- Heinzel, S., Schulte, S., Onken, J., Duong, Q.-L., Riemer, T.G., Heinz, A., Kathmann, N., Rapp, M., 2013. Working memory training improvements and gains in non-trained cognitive tasks in young and older adults. *Neuropsychol. Dev. Cogn. B Aging Neuropsychol. Cogn.* 21, 146–173. <https://doi.org/10.1080/13825585.2013.790338>.
- Heinzel, S., Lorenz, R.C., Pelz, P., Heinz, A., Walter, H., Kathmann, N., Rapp, M.A., Stelzel, C., 2016. Neural correlates of training and transfer effects in working memory in older adults. *Neuroimage* 134, 236–249. <https://doi.org/10.1016/j.neuroimage.2016.03.068>.
- Henmi, M., Copas, J.B., 2010. Confidence intervals for random effects meta-analysis and robustness to publication bias. *Stat. Med.* 29, 2969–2983. <https://doi.org/10.1002/sim.4029>.
- Higgins, J.P.T., Altman, D.G., 2008. Assessing risk of bias in included studies. *Cochrane Handbook for Systematic Reviews of Interventions: Cochrane Book Series.* John Wiley & Sons, Ltd, Chichester, UK, pp. 187–241. <https://doi.org/10.1002/9780470712184.ch8>.
- Higgins, J.P.T., Green, S., 2008. *Cochrane Handbook for Systematic Reviews of Interventions: Cochrane Book Series.* Cochrane Handbook for Systematic Reviews of Interventions: Cochrane Book Series. <https://doi.org/10.1002/9780470712184>.
- Higgins, J.P.T., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analyses. *BMJ Br. Med. J.* 327, 557–560. <https://doi.org/10.1136/bmj.327.7414.557>.
- Huitfeldt, B., Danielson, L., Ebbutt, A., Schmidt, K., 2001. Choice of control in clinical trials - Issues and implications of ICH-E10. *Drug Inf. J.* 35, 1147–1156. <https://doi.org/10.1177/0959286150103500411>.
- ICH Harmonised Tripartite Guideline, 2000. ICH - E10 - Choice of Control Group and Related Issues in Clinical Trials, ICH Harmonised Tripartite Guideline. <https://doi.org/10.1016/j.jaad.2006.09.029>.
- Jaeggi, S.M., Buschkuhl, M., Jonides, J., Perrig, W.J., 2008. Improving fluid intelligence with training on working memory. *Proc. Natl. Acad. Sci.* 105, 6829–6833. <https://doi.org/10.1073/pnas.0801268105>.
- Jaeggi, S.M., Buschkuhl, M., Shah, P., Jonides, J., 2014. The role of individual differences in cognitive training and transfer. *Mem. Cognit.* 42, 464–480. <https://doi.org/10.3758/s13421-013-0364-z>.
- Jolles, D.D., Crone, E.A., 2012. Training the developing brain: a neurocognitive perspective. *Front. Hum. Neurosci.* 6. <https://doi.org/10.3389/fnhum.2012.00076>.
- Just, M.A., Carpenter, P.A., 1992. A capacity theory of comprehension: individual differences in working memory. *Psychol. Rev.* 99, 122–149. <https://doi.org/10.1037/0033-295X.99.1.122>.
- Karbach, J., Verhaeghen, P., 2014. Making working memory work: A meta-analysis of executive-control and working memory training in older adults. *Psychol. Sci.* 25, 2027–2037. <https://doi.org/10.1177/0956797614548725>.
- Kawagoe, T., Suzuki, M., Nishiguchi, S., Abe, N., Otsuka, Y., Nakai, R., Yamada, M., Yoshikawa, S., Sekiyama, K., 2015. Brain activation during visual working memory correlates with behavioral mobility performance in older adults. *Front. Aging Neurosci.* 7, 186. <https://doi.org/10.3389/fnagi.2015.00186>.
- Kelly, M.E., Loughrey, D., Lawlor, B.A., Robertson, I.H., Walsh, C., Brennan, S., 2014. The impact of cognitive training and mental stimulation on cognitive and everyday functioning of healthy older adults: a systematic review and meta-analysis. *Ageing Res. Rev.* <https://doi.org/10.1016/j.arr.2014.02.004>.
- Kemper, S., Herman, R.E., Liu, C.J., 2004. Sentence production by young and older adults in controlled contexts. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 59, P220–4. <https://doi.org/10.1093/geronb/59.5.P220>.
- Ko, P.C., Duda, B., Hussey, E., Mason, E., Molitor, R.J., Woodman, G.F., Ally, B.A., 2014. Understanding age-related reductions in visual working memory capacity: examining the stages of change detection. *Attention Percept. Psychophys.* <https://doi.org/10.3758/s13414-013-0585-z>.
- Laguna, K., Babcock, R.L., 1997. Computer anxiety in young and older adults:

- implications for human-computer interactions in older populations. *Comput. Human Behav.* [https://doi.org/10.1016/S0747-5632\(97\)00012-5](https://doi.org/10.1016/S0747-5632(97)00012-5).
- Lampit, A., Hallock, H., Valenzuela, M., 2014. Computerized cognitive training in cognitively healthy older adults: a systematic review and meta-analysis of effect modifiers. *PLoS Med.* 11. <https://doi.org/10.1371/journal.pmed.1001756>.
- Landi, D., Rossini, P.M., 2010. Cerebral restorative plasticity from normal ageing to brain diseases: a "never ending story". *Restor. Neurol. Neurosci.* <https://doi.org/10.3233/RNN-2010-0538>.
- Landis, J.R., Koch, G.G., 1977. The measurement of observer agreement for categorical data data for categorical of observer agreement the measurement. *Biometrics* 33, 159–174. <https://doi.org/10.2307/2529310>.
- Lange, S., Süß, H.M., 2015. Experimental evaluation of near- and far-transfer effects of an adaptive multicomponent working memory training. *Appl. Cogn. Psychol.* 29, 502–514. <https://doi.org/10.1002/acp.3126>.
- Li, S.-C., Schmiedek, F., Huxhold, O., Röcke, C., Smith, J., Lindenberger, U., 2008. Working memory plasticity in old age: practice gain, transfer, and maintenance. *Psychol. Aging* 23, 731–742. <https://doi.org/10.1037/a0014343>.
- Lilienthal, L., Tamez, E., Shelton, J.T., Myerson, J., Hale, S., 2013. Dual n-back training increases the capacity of the focus of attention. *Psychon. Bull. Rev.* 20, 135–141. <https://doi.org/10.3758/s13423-012-0335-6>.
- Loosli, S.V., Falquez, R., Unterrainer, J.M., Weiller, C., Rahm, B., Kaller, C.P., 2016. Training of resistance to proactive interference and working memory in older adults: a randomized double-blind study. *Int. Psychogeriatrics.* <https://doi.org/10.1017/S1041610215001519>.
- Lubitz, A.F., Niedeggen, M., Feser, M., 2017. Aging and working memory performance: electrophysiological correlates of high and low performing elderly. *Neuropsychologia* 106, 42–51. <https://doi.org/10.1016/j.neuropsychologia.2017.09.002>.
- Lustig, C., Shah, P., Seidler, R., Reuter-Lorenz, P.A., 2009. Aging, training, and the brain: a review and future directions. *Neuropsychol. Rev.* <https://doi.org/10.1007/s11065-009-9119-9>.
- Mansur-Alves, M., Silva, R.S., 2017. Treinar memória de trabalho promove mudanças em inteligência fluida? Temas em Psicol. 25, 787–807. <https://doi.org/10.9788/TP2017.2-19Pt>.
- Melby-Lervåg, M., Hulme, C., 2013. Is working memory training effective? A meta-analytic review. *Dev. Psychol.* 49, 270–291. <https://doi.org/10.1037/a0028228>.
- Melby-Lervåg, M., Hulme, C., 2016. There is no convincing evidence that working memory training is effective: a reply to Au et al. (2014) and Karch and Verhaeghen (2014). *Psychon. Bull. Rev.* 23, 324–330. <https://doi.org/10.3758/s13423-015-0862-z>.
- Melby-Lervåg, M., Redick, T.S., Hulme, C., 2016. Working memory training does not improve performance on measures of intelligence or other measures of "Far transfer": evidence from a meta-analytic review. *Perspect. Psychol. Sci.* 11, 512–534. <https://doi.org/10.3837/tis.0000.00.000>.
- Mlinarić, A., Horvat, M., Smolčić, V.Š., 2017. Dealing with the positive publication bias: why you should really publish your negative results. *Biochem. Medica.* <https://doi.org/10.11613/BM.2017.030201>.
- Moeyaert, M., Ugille, M., Natasha Beretvas, S., Ferron, J., Bunuan, R., Van den Noortgate, W., 2017. Methods for dealing with multiple outcomes in meta-analysis: a comparison between averaging effect sizes, robust variance estimation and multilevel meta-analysis. *Int. J. Soc. Res. Methodol.* 20, 559–572. <https://doi.org/10.1080/13645579.2016.1252189>.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G.D., Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G.D., 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 8 <https://doi.org/10.1136/bmj.b2535>.
- Morris, S.B., 2008. Estimating effect sizes from pretest-posttest-control group designs. *Organ. Res. Methods* 11, 364–386. <https://doi.org/10.1177/1094428106291059>.
- Morrison, A.B., Chein, J.M., 2011. Does working memory training work? The promise and challenges of enhancing cognition by training working memory. *Psychon. Bull. Rev.* 18, 46–60. <https://doi.org/10.3758/s13423-010-0034-0>.
- Neely, A.S., Nyberg, L., 2015. Working Memory Training in Late Adulthood: a Behavioral and Brain perspective, Working Memory and Ageing. *Psychology Press* <https://doi.org/10.4324/9781315879840-10>.
- Nittrouer, S., Lowenstein, J.H., Wucnich, T., Moberly, A.C., 2016. Verbal working memory in older adults: the roles of phonological capacities and processing speed. *J. Speech Lang. Hear. Res.* [https://doi.org/10.1044/2016\\_jslhr-h-15-0404](https://doi.org/10.1044/2016_jslhr-h-15-0404).
- Noack, H., Lövdén, M., Schmiedek, F., Lindenberger, U., 2009. Cognitive plasticity in adulthood and old age: gauging the generality of cognitive intervention effects. *Restor. Neurol. Neurosci.* 27, 435–453. <https://doi.org/10.3233/RNN-2009-0496>.
- Oberauer, K., Süß, H.M., Wilhelm, O., Wittmann, W.W., 2008. Which working memory functions predict intelligence? *Intelligence* 36, 641–652. <https://doi.org/10.1016/j.intell.2008.01.007>.
- Oberauer, K., Lewandowsky, S., Awh, E., Brown, G.D.A., Conway, A., Cowan, N., Donkin, C., Farrell, S., Hitch, G.J., Hurlstone, M.J., Ma, W.J., Morey, C.C., Nee, D.E., Schweppke, J., Vergauwe, E., Ward, G., 2018. Benchmarks for models of short-term and working memory. *Psychol. Bull.* <https://doi.org/10.1037/bul0000153>.
- Olson, I.R., Zhang, J.X., Mitchell, K.J., Johnson, M.K., Bloise, S.M., Higgins, J.A., 2004. Preserved spatial memory over brief intervals in older adults. *Psychol. Aging.* <https://doi.org/10.1037/0882-7974.19.2.310>.
- Oswald, W.D., Gunzelmann, T., Rupprecht, R., Hagen, B., 2006. Differential effects of single versus combined cognitive and physical training with older adults: the SimA study in a 5-year perspective. *Eur. J. Ageing* 3, 179–192. <https://doi.org/10.1007/s10433-006-0035-z>.
- Park, D.C., Reuter-Lorenz, P., 2009. The adaptive brain: aging and neurocognitive scaffolding. *Annu. Rev. Psychol.* 60, 173–196. <https://doi.org/10.1146/annurev.psych.59.103006.093656>.
- Penner, I.-K., Vogt, A., Stöcklin, M., Gschwind, L., Opwis, K., Calabrese, P., 2012. Computerized working memory training in healthy adults: a comparison of two different training schedules. *Neuropsychol. Rehabil.* 22, 716–733. <https://doi.org/10.1080/09602011.2012.686883>.
- Pergher, V., Wittevröngel, B., Tournay, J., Schoenmakers, B., Van Hulle, M.M., 2018. N-back training and transfer effects revealed by behavioral responses and EEG. *Brain Behav.* <https://doi.org/10.1002/brb3.1136>.
- Pustejovsky, J., 2017. Package "clubSandwich" Title Cluster-robust (Sandwich) Variance Estimators with Small-sample Corrections. <https://doi.org/10.1080/07350015.2016.1247004&gt>.
- R Core Team, 2018. R: A Language and Environment for Statistical Computing [WWW Document]. Vienna, Austria. URL. <https://www.r-project.org/>.
- Ragland, J.D., Turetsky, B.I., Gur, R.C., Gunning-Dixon, F., Turner, T., Schroeder, L., Chan, R., Gur, R.E., 2002. Working memory for complex figures: an fMRI comparison of letter and fractal n-back tasks. *Neuropsychology.* <https://doi.org/10.1037/0894-4105.16.3.370>.
- Raz, N., 2005. The Aging Brain Observed in Vivo: Differential Changes and Their Modifiers, Cognitive Neuroscience of Aging: Linking Cognitive and Cerebral Aging Cabeza, R. Nyberg, L. Park, D. <https://doi.org/10.1176/appi.ajp.163.3.560>.
- Richmond, L.L., Morrison, A.B., Chein, J.M., Olson, I.R., 2011. Working memory training and transfer in older adults. *Psychol. Aging* 26, 813–822. <https://doi.org/10.1037/a0023631>.
- Rothwell, P.M., 2006. Factors that can affect the external validity of randomised controlled trials. *PLOS Hub Clin. Trials* 1, e9. <https://doi.org/10.1371/journal.pctr.0010009>.
- Rottschy, C., Langner, R., Dogan, I., Reetz, K., Laird, A.R., Schulz, J.B., Fox, P.T., Eickhoff, S.B., 2012. Modelling neural correlates of working memory: a coordinate-based meta-analysis. *Neuroimage* 60, 830–846. <https://doi.org/10.1016/j.neuroimage.2011.11.050>.
- Salthouse, T.A., 1990. Working memory as a processing resource in cognitive aging. *Dev. Rev.* 10, 101–124. [https://doi.org/10.1016/0273-2297\(90\)90006-P](https://doi.org/10.1016/0273-2297(90)90006-P).
- Salthouse, T.A., 2000. Aging and measures of processing speed. *Biol. Psychol.* [https://doi.org/10.1016/S0301-0511\(00\)00052-1](https://doi.org/10.1016/S0301-0511(00)00052-1).
- Schmiedek, F., Hildebrandt, A., Lövdén, M., Wilhelm, O., Lindenberger, U., 2009. Complex span versus updating tasks of working memory: the gap is not that deep. *J. Exp. Psychol. Learn. Mem. Cogn.* 35, 1089–1096. <https://doi.org/10.1037/a0015730>.
- Schwaighofer, M., Fischer, F., Bühner, M., 2015. Does working memory training transfer? A meta-analysis including training conditions as moderators. *Educ. Psychol.* 50, 138–166. <https://doi.org/10.1080/00461520.2015.1036274>.
- Shakeel, M.K., Goghari, V.M., 2017. Measuring fluid intelligence in healthy older adults. *J. Aging Res.* <https://doi.org/10.1155/2017/8514582>.
- Shing, Y.L., Schmiedek, F., Lövdén, M., Lindenberger, U., 2012. Memory updating practice across 100 days in the COGITO study. *Psychol. Aging.* <https://doi.org/10.1037/a0025568>.
- Shipstead, Z., Redick, T.S., Engle, R.W., 2012. Is working memory training effective? *Psychol. Bull.* 138, 628–654. <https://doi.org/10.1037/a0027473>.
- Simon, S.S., Tusch, E.S., Feng, N.C., Häkansson, K., Mohammed, A.H., Daffner, K.R., 2018. Is computerized working memory training effective in healthy older adults? Evidence from a multi-site, randomized controlled trial. *J. Alzheimers Dis.* <https://doi.org/10.3233/JAD-180455>.
- Stepankova, H., Lukavsky, J., Buschkuhl, M., Kopecek, M., Ripova, D., Jaeggi, S.M., 2014. The malleability of working memory and visuospatial skills: a randomized controlled study in older adults. *Dev. Psychol.* 50, 1049–1059. <https://doi.org/10.1037/a0034913>.
- Sterne, J.A., Egger, M., Moher, D., 2008. Addressing reporting biases. *Cochrane Handbook for Systematic Reviews of Interventions: Cochrane Book Series.* pp. 297–333. <https://doi.org/10.1002/9780470712184.ch10>.
- Sterne, J.A.C., Sutton, A.J., Ioannidis, J.P.A., Terrin, N., Jones, D.R., Lau, J., Carpenter, J., Rücker, G., Harbord, R.M., Schmid, C.H., Tetzlaff, J., Deeks, J.J., Peters, J., Macaskill, P., Schwarzer, G., Duval, S., Altman, D.G., Moher, D., Higgins, J.P.T., 2011. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 343, d4002. <https://doi.org/10.1136/bmj.d4002>.
- Takeuchi, H., Taki, Y., Nouchi, R., Hashizume, H., Sekiguchi, A., Kotozaki, Y., Nakagawa, S., Miyauchi, C.M., Sassa, Y., Kawashima, R., 2013. Effects of working memory training on functional connectivity and cerebral blood flow during rest. *Cortex* 49, 2106–2125. <https://doi.org/10.1016/j.cortex.2012.09.007>.
- Takeuchi, H., Taki, Y., Nouchi, R., Hashizume, H., Sekiguchi, A., Kotozaki, Y., Nakagawa, S., Miyauchi, C.M., Sassa, Y., Kawashima, R., 2014. Working memory training improves emotional states of healthy individuals. *Front. Syst. Neurosci.* 8, 200. <https://doi.org/10.3389/fnsys.2014.00200>.
- Teixeira-Santos, A.A.C., Nafee, T., Sampaio, A., Leite, J., Carvalho, S., 2015. Effects of transcranial direct current stimulation on working memory in healthy older adults: a systematic review. *PPCR* 1 (3), 73–81.
- Tkatch, R., Musich, S., MacLeod, K., Hawkins, K., Yeh, C.S., 2016. Population health management for older adults. *Gerontol. Geriatr. Med.* <https://doi.org/10.1177/2333721416667877>.
- Toril, P., Reales, J.M., Ballesteros, S., 2014. Video game training enhances cognition of older adults: a meta-analytic study. *Psychol. Aging.* <https://doi.org/10.1037/a0037507>.
- United Nations, Department of Economic and Social Affairs, P.D., 2017. World Population Ageing 2017. United Nations 124. <https://doi.org/ST/ESA/SER.A/348>.
- Unsworth, N., Heitz, R.P., Engle, R.W., 2005. Working memory capacity in Hot and cold cognition. *Cognitive Limitations in Aging and Psychopathology.* pp. 19–43. <https://doi.org/10.1017/CBO9780511720413.003>.

- Verhaeghen, P., Marcoen, A., Goossens, L., 1992. Improving memory performance in the aged through mnemonic training: a meta-analytic study. *Psychol. Aging* 7, 242–251. <https://doi.org/10.1037/0882-7974.8.3.338>.
- Viechtbauer, W., 2010. Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.* 36, 1–48. <https://doi.org/10.1103/PhysRevB.91.121108>.
- Viswanathan, M., Patnode, C.D., Berkman, N.D., Bass, E.B., Chang, S., Hartling, L., Murad, M.H., Treadwell, J.R., Kane, R.L., 2008. Assessing the Risk of Bias in Systematic Reviews of Health Care Interventions, Methods Guide for Effectiveness and Comparative Effectiveness Reviews. Agency for Healthcare Research and Quality (US).
- von Bastian, C.C., Eschen, A., 2016. Does working memory training have to be adaptive? *Psychol. Res.* 80, 181–194. <https://doi.org/10.1007/s00426-015-0655-z>.
- von Bastian, C.C., Oberauer, K., 2013. Effects and mechanisms of working memory training: a review. *Psychol. Res.* 1–18. <https://doi.org/10.1007/s00426-013-0524-6>.
- von Bastian, C.C., Langer, N., Jäncke, L., Oberauer, K., 2013. Effects of working memory training in young and old adults. *Mem. Cognit.* 41, 611–624. <https://doi.org/10.3758/s13421-012-0280-7>.
- Wayne, R.V., Hamilton, C., Huyck, J.J., Johnsrude, I.S., 2016. Working memory training and speech in noise comprehension in older adults. *Front. Aging Neurosci.* <https://doi.org/10.3389/fnagi.2016.00049>.
- Weicker, J., Villringer, A., Thöne-Otto, A., 2016. Can impaired working memory functioning be improved by training? A meta-analysis with a special focus on brain injured patients. *Neuropsychology* 30, 190–212. <https://doi.org/10.1037/neu0000227>.
- Weicker, J., Hudl, N., Frisch, S., Lepsien, J., Mueller, K., Villringer, A., Thöne-Otto, A., 2018. WOME: Theory-Based Working Memory Training — A Placebo-Controlled, Double-Blind Evaluation in Older Adults. *Front. Aging Neurosci.* 10, 247. <https://doi.org/10.3389/fnagi.2018.00247>.
- West, R., 1999. Visual distraction, working memory, and aging. *Mem. Cogn.* <https://doi.org/10.3758/BF03201235>.
- Wickstrom, G., Bendix, T., 2000. The “Hawthorne effect” - what did the original Hawthorne studies actually show? *Scand. J. Work. Environ. Heal.* <https://doi.org/10.5271/sjweh.555>.
- Xin, Z., Lai, Z.R., Li, F., Maes, J.H.R., 2014. Near- and far-transfer effects of working memory updating training in elderly adults. *Appl. Cogn. Psychol.* 28, 403–408. <https://doi.org/10.1002/acp.3011>.
- Zajac-Lamparska, L., Trempala, J., 2016. Effects of working memory and attentional control training and their transfer onto fluid intelligence in early and late adulthood. *Heal. Psychol. Rep.* <https://doi.org/10.5114/hpr.2016.56846>.
- Zhou, X., Ye, Y., Tang, G., Wu, F., 2017. Small-study effects” in meta-analysis should not be ignored. *J. Crit. Care* 39, 283–284. <https://doi.org/10.1016/j.jcrc.2017.01.013>.
- Zinke, K., Zeintl, M., Eschen, A., Herzog, C., Kliegel, M., 2011. Potentials and limits of plasticity induced by working memory training in old-old age. *Gerontology* 58, 79–87. <https://doi.org/10.1159/000324240>.
- Zinke, K., Zeintl, M., Rose, N.S., Putzmann, J., Pydde, A., Kliegel, M., 2014. Working memory training and transfer in older adults: effects of age, baseline performance, and training gains. *Dev. Psychol.* <https://doi.org/10.1037/a0032982>.